

# Package ‘DNAseqtest’

October 12, 2022

**Type** Package

**Title** Generating and Testing DNA Sequences

**Version** 1.0

**Date** 2016-03-26

**Author** Faisal Ababneh, John Robinson, Lars S Jermiin and Hasinur Rahaman Khan

**Maintainer** Hasinur Rahaman Khan <hasinurkhan@gmail.com>

**Description** Generates DNA sequences based on Markov model techniques for matched sequences. This can be generalized to several sequences. The sequences (taxa) are then arranged in an evolutionary tree (phylogenetic tree) depicting how taxa diverge from their common ancestors. This gives the tests and estimation methods for the parameters of different models. Standard phylogenetic methods assume stationarity, homogeneity and reversibility for the Markov processes, and often impose further restrictions on the parameters.

**License** GPL-2

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2016-03-24 23:32:09

## R topics documented:

DNAseqtest-package . . . . .	2
artomat . . . . .	3
Distance . . . . .	4
Fmatrix . . . . .	5
gn . . . . .	6
gn2 . . . . .	7
gn3sim . . . . .	8
likelihood . . . . .	9
Ntml . . . . .	10
Pt . . . . .	11
simapp . . . . .	12
simemb . . . . .	13
Smatrix . . . . .	14
TEST2 . . . . .	15
TEST3 . . . . .	16

---

 DNaseqtest-package      *Generating and Testing DNA Sequences*


---

### Description

Generates DNA sequences based on Markov model techniques for matched sequences. This can be generalized to several sequences. The sequences (taxa) are then arranged in an evolutionary tree (phylogenetic tree) depicting how taxa diverge from their common ancestors. This gives the tests and estimation methods for the parameters of different models. Standard phylogenetic methods assume stationarity, homogeneity and reversibility for the Markov processes, and often impose further restrictions on the parameters.

### Details

Package: DNaseqtest  
 Type: Package  
 Version: 1.0  
 Date: 2016-03-26  
 License: GPL-2

### Author(s)

Faisal Ababneh, John Robinson, Lars S Jermiin and Hasinur Rahaman Khan Maintainer: Hasinur Rahaman Khan <hasinurkhan@gmail.com>

### References

Lars Sommer Jermiin, Vivek Jayaswal, Faisal Ababneh, John Robinson (2008). Phylogenetic model evaluation. *Bioinformatics*, Volume 452 of the series *Methods in Molecular Biology*, 331-364.

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

Faisal Ababneh, Lars S Jermiin, John Robinson (2006). Generation of the Exact Distribution and Simulation of Matched Nucleotide Sequences on a Phylogenetic Tree. *Journal of mathematical modelling and algorithms*, 5(3), 291-308.

### Examples

```
#To generate a 4^5 gene array
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3),4,2)
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.35,.79,.01,.93,.47), 3,.1,.5,.8)
gn.sec<-gn(theta, merge2)
```

gn.sec

---

artomat

*Transforming  $4^K$  Array to  $m \times K$  Matrix*

---

### Description

This function transfers any array to a matrix.

### Usage

```
artomat(fobs)
```

### Arguments

fobs            a  $4^K$  array, containing the observed divergent frequencies for K matched sequences

### Details

This function transfers any  $4^K$  array containing the observed divergent frequencies of K matched sequences to an  $m \times K$  matrix, where m is the sum of the frequencies in the  $4^K$  observed divergence array.

### Value

An  $m \times K$  matrix

### References

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

### See Also

gn2, gn, Fmatrix

---

 Distance

*Paralinear Distances*


---

### Description

This function calculates the paralinear distance between K matched DNA sequences.

### Usage

```
Distance(F4)
```

### Arguments

F4                    a  $4^K$  array containing the joint distribution array  $F(t)$  or the observed array N

### Details

This function calculates the paralinear distances between K matched DNA sequences, depending on the joint distribution array for these K sequences or on the observed divergence array N.

### Value

A K x K symmetric matrix distances between the K sequences

### References

Faisal Ababneh, Lars S Jermin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

### See Also

gn2, gn, Fmatrix, Ntml

### Examples

```
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.35,.79,.01,.93,.47), 3,.1,.5,.8)
F1<-gn(theta,merge2)
dn<-Distance(F1)
dn
```

**Description**

This function calculates the joint distribution function for two edge tree.

**Usage**

```
Fmatrix(t1, t2, f0, Sx2, Sy2, Pix, Piy)
```

**Arguments**

t1	represents the length from the tree root to the first node
t2	represents the length from the tree root to the second node
f0	the initial distribution for the four nucleotides
Sx2	a 4 x 4 symmetric matrix related to the first edge
Sy2	a 4 x 4 symmetric matrix related to the second edge
Pix	a diagonal matrix for the stationary distribution of the first edge
Piy	a diagonal matrix for the stationary distribution of the second edge

**Details**

This function calculates the joint distribution function for a two edge tree with different edge lengths, stationary distributions and differentS matrices.

**Value**

A 4 x 4 matrix containing the joint edges

**References**

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

gn, Smatrix

**Examples**

```
f0<-c(.25,.25,.25,.25)
Pi1<-diag(c(.2,.2,.2,.4))
Pi2<-diag(c(.1,.1,.1,.7))
S1<-Smatrix(c(.2,.2,.2,.2,.2,.2),diag(Pi1))
S2<-Smatrix(c(.3,.3,.3,.3,.3,.3),diag(Pi2))
fm<-Fmatrix(1,.5,f0,S1,S2,Pi1,Pi2)
fm
```

gn

*Joint Distribution for K Matched Sequences***Description**

This function calculates the joint distribution array for K matched sequences.

**Usage**

```
gn(theta, merge2)
```

**Arguments**

theta	a vector of variables containing the following parameters in this order–1. the first three parameters from $\pi_X$ vector, 2. the first three parameters from $\pi_Y$ vector, 3. the first three parameters from $f_0$ vector, 4. the six off-diagonal free parameters in the S matrix, 5. a scalar $\rho$ , 6. a vector of lengths containing K-2 values
merge2	(K-1) x 2 matrix describing the tree topology

**Details**

This function calculates the joint distribution array for a tree with K matched sequences. it uses the following functions– Pt, Fmatrix and Smatrix.

**Value**

A  $4^K$  array containing the joint distribution for the K edges

**References**

Lars Sommer Jermiin, Vivek Jayaswal, Faisal Ababneh, John Robinson (2008). Phylogenetic model evaluation. *Bioinformatics*, Volume 452 of the series *Methods in Molecular Biology*, 331-364.

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

Fmatrix, Pt, Smatrix

**Examples**

```
#To generate a 4^5 gene array
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3),4,2)
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.35,.79,.01,.93,.47), 3,.1,.5,.8)
gn.sec<-gn(theta, merge2)
gn.sec
```

gn2

*Joint Distribution for K Matched Sequences (2)***Description**

This function calculates the joint distribution array for K matched sequences (second option).

**Usage**

```
gn2(theta, merge2)
```

**Arguments**

theta	a vector of variables containing the following parameters in this order–1. the first three parameters from $\pi_X$ vector, 2. the first three parameters from $\pi_Y$ vector, 3. the first three parameters from $f_0$ vector, 4. the six off-diagonal free parameters in the S matrix, 5. a scalar $\rho$ , 6. a vector of lengths containing K-2 values
merge2	(K-1) x 2 matrix describing the tree topology

**Details**

This function calculates the joint distribution array for a tree with K matched sequences. it uses the following functions– Pt, Fmatrix and Smatrix.

**Value**

A  $4^K$  array containing the joint distribution for the K edges

**References**

Lars Sommer Jermiin, Vivek Jayaswal, Faisal Ababneh, John Robinson (2008). Phylogenetic model evaluation. Bioinformatics, Volume 452 of the series Methods in Molecular Biology, 331-364.

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. Bioinformatics, 22(10), 1225-1231.

**See Also**

Fmatrix, Pt, Smatrix

**Examples**

```
#To generate a 4^5 gene array
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
rho2<-matrix(c(.3,.5,.3,.2,.3,.5,.8,2.7),4,2)
theta<-c(rep(.25,3), rep(.25,3),rep(.25,3), c(.2,.35,.79,.01,.93,.47),rho2)
gn2<-gn2(theta, merge2)
gn2
```

gn3sim

---

*Generating Random DNA Samples Using the Rambaut and Grassly Method*

---

**Description**

This function generates random DNA samples using Rambaut and Grassly method.

**Usage**

```
gn3sim(theta, seqLength, merge2)
```

**Arguments**

theta	a vector of variables containing the following parameters in this order–1. the first three parameters from $\pi_X$ vector, 2. the first three parameters from $\pi_Y$ vector, 3. the first three parameters from $f_0$ vector, 4. the six off-diagonal free parameters in the S matrix, 5. a scalar $\rho$ , 6. a vector of lengths containing K-2 values
seqLength	the length of sequences we need to generate
merge2	(K-1) x 2 matrix describing the tree topology

**Details**

This function generates a  $4^K$  DNA array using Rambaut and Grassly, (1997) method. It depends on a set of variables theta, the sequence length and a merge matrix describing the tree topology.

**Value**

A n x K observed divergence matrix

**References**

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.



**See Also**

Ntml, simapp, simemb, gn, gn2, Fmatrix

**Examples**

```
# This will give 4^5 observed divergence array
theta<-(c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2, .35, .79, .01, .93, .47),
3, .1, .5, .8))
n<-1000
merge2<-matrix(c(-1, -4, -3, 2, -2, -5, 1, 3), 4, 2)
gn3<-gn3sim(theta, n, merge2)
gn3
```

---

likelihood

*Negative Log Likelihood Ratio*

---

**Description**

This function calculates log likelihood ratio value.

**Usage**

```
likelihood(thetast, fobs, merge2)
```

**Arguments**

thetast	a starting values for the parameter we need to estimate
fobs	the $4^K$ joint distribution array for K edge tree
merge2	a (K-1) x 2 matrix describing the tree topology

**Details**

This function calculates the log likelihood ratio value for F(t). It needs a vector of starting values for the parameters estimate,  $4^K$  observed divergence array and merge matrix describing the tree topology.

**Value**

The value of the log likelihood ratio

**References**

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

gn, gn2

**Examples**

```
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.35,.79,.01,.93,.47),3,.1,.5,.8)
F1<-gn(theta, merge2)
lh<-likelihood(theta, F1, merge2)
lh
```

---

Ntml

*Generating Samples from a Multinomial Distribution*

---

**Description**

Generating random DNA samples from a multinomial distribution.

**Usage**

```
Ntml(N, Ft)
```

**Arguments**

N	sample size
Ft	a $4^K$ array, containing the joint distribution probabilities for K matched sequences.

**Details**

This function generates a  $4^K$  DNA array from a multinomial distribution. It depends on the sample size we need to generate and the  $4^K$  joint distribution array of K matched sequences.

**Value**

A  $4^K$  observed divergence array

**References**

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

simemb, simapp, gn3sim, gn, gn2, Fmatrix

## Examples

```
#This will give a 4^K observed divergence array
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.35,.79,.01,.93,.47),
3,.1,.5,.8)
F1<-gn(theta,merge2)
Nt<-Ntml(1000, F1)
Nt
```

---

Pt *Transition Probability Function*

---

## Description

This function calculates the transition probability function for a process during a period of time.

## Usage

```
Pt(S, Pi, t)
```

## Arguments

S	a 4 x 4 symmetric matrix
Pi	a diagonal matrix containing the stationary distribution for the process
t	a period of time describing the length of the process

## Details

This function needs the 4 x 4 symmetric matrix S,  $\Pi$  and the process length t in order to find the transition probability over that process, where  $P_{ij}(t)$  is the probability that the *i*th nucleotide changes to the *j*-th nucleotide during the period of t.

## Value

A 4 x 4 matrix containing the transition probabilities for a process.

## References

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

## See Also

Smatrix

**Examples**

```
Pi<-diag(c(.1,.1,.1,.7))
S<-Smatrix(c(.3,.3,.3,.3,.3,.3),diag(Pi))
t<-1
p<-Pt(S, Pi, t)
p
```

---

simapp

*Generating Random DNA Samples Using Approximation Method*


---

**Description**

This function generates random DNA samples using an approximation method

**Usage**

```
simapp(theta, seqLength, merge1)
```

**Arguments**

theta	a vector of variables containing the following parameters in this order—1. the first three parameters from $\pi_X$ vector, 2. the first three parameters from $\pi_Y$ vector, 3. the first three parameters from $f_0$ vector, 4. the six off-diagonal free parameters in the S matrix, 5. a scalar $\rho$ , 6. a vector of lengths containing K-2 values
seqLength	the length of sequences we need to generate
merge1	(K-1) x 2 matrix describing the tree topology

**Details**

This function generates a  $4^K$  DNA array using an approximation method. It depends on a set of variables theta, the sequence length and a merge matrix describing the tree topology.

**Value**

A n x K observed divergence matrix

**References**

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

Ntml, simemb, gn3sim, gn, gn2, Fmatrix

**Examples**

```
# This will give 4^5 observed divergence array
theta<-(c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.2,.2,.2,.2,.2),
3,.1,.5,.8))
n<-1000
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
sa<-simapp(theta, n, merge2)
sa
```

simemb

*Generating Random DNA Samples Using Embedded Markov Chain***Description**

This function generates random DNA samples using embedded chain.

**Usage**

```
simemb(theta, seqLength, merge2)
```

**Arguments**

theta	a vector of variables containing the following parameters in this order–1. the first three parameters from $\pi_X$ vector, 2. the first three parameters from $\pi_Y$ vector, 3. the first three parameters from $f_0$ vector, 4. the six off-diagonal free parameters in the S matrix, 5. a scalar $\rho$ , 6. a vector of lengths containing $K-2$ values
seqLength	the length of sequences we need to generate
merge2	$(K-1) \times 2$ matrix describing the tree topology

**Details**

This function generates  $4^K$  DNA array using embedded Markov chain. It depends on a set of variables theta, the sequence length and a merge matrix describing the tree topology.

**Value**

A  $n \times K$  observed divergence matrix

**References**

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

Ntml, simapp, gn3sim, gn, gn2, Fmatrix

**Examples**

```
# This will give 4^5 observed divergence array
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2, .35, .79, .01, .93, .47),
3, .1, .5, .8))
n<-1000
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
sm<-simemb(theta, n, merge2)
sm
```

Smatrix

*Symmetric Matrix S***Description**

This function calculates the symmetric matrix S.

**Usage**

```
Smatrix(s, pi)
```

**Arguments**

s a vector of variables containing the six free parameters in the S matrix  
pi a vector giving the stationary probabilities for the four nucleotides A, C, G and T

**Details**

This function calculates the matrix S, which we used to calculate the rate matrix R.

**Value**

A 4 x 4 symmetric matrix

**References**

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

Pt, Fmatrix, gn ,gn2

**Examples**

```
s<-c(.1, .2, .3, .4, .5, .6)
pi<-c(.1, .1, .1, .7)
sm<-Smatrix(s, pi)
sm
```

---

TEST2

*Test for Symmetry of Matched DNA Sequences*

---

### Description

This function tests for symmetry between all the pairs of K matched DNA sequences.

### Usage

TEST2(f)

### Arguments

f                    a  $4^K$  array containing the observed divergence array N

### Details

This function calculates Bowker's test for symmetry, Stuart's test for marginal symmetry and the test for internal symmetry. It depends on the  $4^K$  observed divergence array N.

### Value

A list of three lower triangle matrices

first	the lower triangle of the matrix contains (K-1) x (K-1) values shows Bowker's test between all the possible pairs of the K sequences
second	the lower triangle of the matrix contains (K-1) x (K-1) values shows Stuart's test between all the possible pairs of the K sequences
third	the lower triangle of the matrix contains (K-1) x (K-1) values shows the internal test between all the possible pairs of the K sequences

### References

Lars Sommer Jermiin, Vivek Jayaswal, Faisal Ababneh, John Robinson (2008). Phylogenetic model evaluation. *Bioinformatics*, Volume 452 of the series *Methods in Molecular Biology*, 331-364.

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

### See Also

Ntml, simapp, simemb, TEST3

**Examples**

```
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.35,.79,.01,.93,.47),
3,.1,.5,.8)
F1<-gn(theta,merge2)
N1<-Ntml(1000,F1)
t2<-TEST2(N1)
t2
```

---

TEST3

---

*Overall Test for Marginal Symmetry*


---

**Description**

This function tests for symmetry between  $K$  matched DNA sequences.

**Usage**

```
TEST3(Farray)
```

**Arguments**

Farray            a  $4^K$  array containing the observed divergence array  $N$

**Details**

This function calculates overall test for marginal symmetry. It depends on the  $4^K$  observed divergence array  $N$ .

**Value**

A single value gives the overall test for marginal symmetry between  $K$  matched sequences

**References**

Lars Sommer Jermiin, Vivek Jayaswal, Faisal Ababneh, John Robinson (2008). Phylogenetic model evaluation. *Bioinformatics*, Volume 452 of the series *Methods in Molecular Biology*, 331-364.

Faisal Ababneh, Lars S Jermiin, Chunsheng Ma, John Robinson (2006). Matched-pairs tests of homogeneity with applications to homologous nucleotide sequences. *Bioinformatics*, 22(10), 1225-1231.

**See Also**

Ntml, simapp, simemb, TEST2



**Examples**

```
merge2<-matrix(c(-1,-4,-3,2,-2,-5,1,3), 4, 2)
theta<-c(rep(.25,3), rep(.25,3), rep(.25,3), c(.2,.35,.79,.01,.93,.47),
3,.1,.5,.8)
F1<-gn(theta,merge2)
N1<-Ntml(1000,F1)
t3<-TEST3(N1)
t3
```

# Index

- \* **Array**
    - artomat, [3](#)
  - \* **Distance between sequences**
    - Distance, [4](#)
  - \* **Generate sequence**
    - gn3sim, [8](#)
    - simapp, [12](#)
    - simemb, [13](#)
  - \* **Generated sample**
    - Ntml, [10](#)
  - \* **Joint distribution**
    - Fmatrix, [5](#)
    - gn, [6](#)
    - gn2, [7](#)
  - \* **Likelihood**
    - likelihood, [9](#)
  - \* **Sequence test**
    - DNaseqtest-package, [2](#)
    - TEST2, [15](#)
    - TEST3, [16](#)
  - \* **Symmetry matrix**
    - Smatrix, [14](#)
  - \* **Transition probability**
    - Pt, [11](#)
- artomat, [3](#)
- Distance, [4](#)
- DNaseqtest (DNaseqtest-package), [2](#)
- DNaseqtest-package, [2](#)
- Fmatrix, [5](#)
- gn, [6](#)
- gn2, [7](#)
- gn3sim, [8](#)
- likelihood, [9](#)
- Ntml, [10](#)
- Pt, [11](#)
- simapp, [12](#)
- simemb, [13](#)
- Smatrix, [14](#)
- TEST2, [15](#)
- TEST3, [16](#)